The Cochrane Library - 1998 Issue 4

Search
Clear
Print+Save
Setup
Handbook
Glossary
Help
Exit

- The Cochrane Database of Systematic Reviews (0 hits, 919 total)
- Database of Abstracts of Reviews of Effectiveness (0 hits, 1820 total)
- The Cochrane Controlled Trials Register (CENTRAL/CCTR) (0 hits, 218356 total)
- The Cochrane Review Methodology Database (0 hits, 836 total)
- About the Cochrane Collaboration (0 hits, 92 total)
- Other sources of information (0 hits, 487 total)

Comments/Criticisms | Back | Find | Outline



THE COCHRANE LIBRARY

1998 Issue 4

ISSN 1464-780X

The best single source of reliable evidence about the effects of health care

Stop Press - Release notes - Getting Started

About The Cochrane Library - Subscription Information - Technical Support

Tasks

Off-line

Search
Clear
Print+Save
Setup
Handbook
Glossary
Help
Exit

Search

Simple Search | Advanced Search | MeSH | Save and Load

"BLOOD PRESSURE" AND STROKE*

Search

Instructions

Enter the term you are looking for in the window above, combining words with the boolean AND, OR or NOT operators. Use the keyword NEAR to search for words near each other, or enclose phrases in quotes to search for words next to each other.

Click the "Search" button (above) or press RETURN to have the number of hits your search returned displayed in the index window, then double-click the database that contains the hits you want to see. The number of hits is printed in red after each database entry, together with the total number of items in that database.

Click New Items or Updated Items (below) to find items new or updated in this issue

For more sophisticated search options, choose "Advanced Search" from this page.

New items only

Updated Items Only

Show the results of the search in the Index window

Help

Cancel


Search
Simple Search
Advanced Search
MeSH
Save and Load
BREAST
Thesaurus
BREAST
ADENOSIS-OF-BREAST see FIBROCYSTIC-DISEASE-OF-BREAST
BREAST
BREAST-DISEASES
BREAST-FEEDING
BREAST-IMPLANTATION
BREAST-IMPLANTS
BREAST-NEOPLASMS
BREAST-NEOPLASMS-MALE
BREAST-SELF-EXAMINATION
BREAST-CYSTS see FIBROCYSTIC-DISEASE-OF-BREAST
BREAST-DYSPLASIA see FIBROCYSTIC-DISEASE-OF-BREAST
BREAST-PROSTHESIS-INTERNAL see BREAST-IMPLANTS
CYSTIC-DISEASE-OF-BREAST see FIBROCYSTIC-DISEASE-OF-BREAST
Single term
Explode in selected tree
Explode in all trees
Choose
Search
Show the results of the search in the Index window
Help
Cancel


Search
Clear
Print+Save
Setup
Handbook
Glossary
Help
Exit
CSF tapping after IVH
Compression for venous leg ulcers
DVT prevention heparin + physical for hip fracture
Doppler in high risk pregnancies
Electrostimulation in placental insufficiency
Hormone therapy for impaired fetal growth
Comments/Criticisms
Back
Find
Outline
Doppler in high risk pregnancies
Doppler ultrasound in high risk pregnancies
Neilson JP, Alfirevic Z
Date of most recent substantive amendment : 23 February 1996
Objectives: To assess whether the use of Doppler ultrasound in high risk pregnancies to investigate flow velocity waveforms in the umbilical artery (+/- uteroplacental arteries) improves obstetric care and fetal outcome.
Search strategy: The register of clinical trials maintained and updated by the Cochrane Pregnancy and Childbirth Group.
Selection criteria: We have identified 20 randomised trials of the use of Doppler ultrasound to investigate umbilical artery waveforms during pregnancy. Eleven of these trials fulfilled prestated criteria for inclusion in this review; in particular, they assessed only women with 'high risk' pregnancies (variously defined). Trial quality
Off-line


The Cochrane Library - 1998 Issue 2
Search
Clear
Print+Save
Setup
Handbook
Glossary
Help
Exit
GnRHa/hMG/FSH for PCOS
HIV infection in pregnancy
Outreach visits
New Ovarian ablation in early breast cancer
New Selegiline in Alzheimer's disease
Spironolactone for hirsutism and/or acne
Comments/Criticisms
Back
Find
Outline
are nearing the menopause,[8,9] ovarian ablation may produce a much greater change in hormonal status in women who do not receive chemotherapy than in those who do. Analyses subdivided by whether or not women in both treatment groups were to receive routine cytotoxic chemotherapy are shown in the figure. The proportional improvement in recurrence-free survival among women aged under 50 at entry was 25% (SD 7) in the absence of chemotherapy but only 10% (9) in its presence (see figure). If real, the difference between these two results could be
Find
Find What:
menopause
Find Next
Cancel
Match Whole Word Only
Match Case
Direction
Up
Down
with ovarian ablation producing a proportional improvement in recurrence-free survival of 27% (7) in the absence of chemotherapy.
POSSIBLE INFLUENCE OF NODAL STATUS IN TRIALS WITHOUT CYTOTOXIC CHEMOTHERAPY
Off-line

Doppler ultrasound in high risk pregnancies

Neilson JP, Alfirevic Z

Date of most recent substantive amendment : 23 February 1996

Objectives: *To assess whether the use of Doppler ultrasound in high risk pregnancies to investigate flow velocity waveforms in the umbilical artery (+/- uteroplacental arteries) improves obstetric care and fetal outcome.*

Search strategy: *The register of clinical trials maintained and updated by the Cochrane Pregnancy and Childbirth Group.*

Selection criteria: *We have identified 20 randomised trials of the use of Doppler ultrasound to investigate umbilical artery waveforms during pregnancy. Eleven of these trials fulfilled prestated criteria for inclusion in this review; in particular, they assessed only women with 'high risk' pregnancies (variously defined). Trial quality was assessed and data were extracted according to the methods employed by the Cochrane Pregnancy and Childbirth Group.*

Data collection and analysis: *Data have been extracted by both authors from published work, complemented, where necessary, by obtaining unpublished data from primary researchers.*

Main results: *The use of Doppler ultrasound in high risk pregnancy (mainly pregnancy associated hypertension and suspected intrauterine growth restriction)*

ID: CN-00164989 - CCTR

TI: **A prospective randomized clinical trial of perioperative treatment with octreotide in pancreas transplantation.**

AU: Benedetti E; Coady NT; Asolati M; Dunn T; Stormoen BM; Bartholomew AM; Vasquez EM; Pollak R

SO: *Am-J-Surg*
YR: 1998
VL: 175
PG: 14-7

AB: BACKGROUND: Technical failures continue to plague clinical pancreas transplantation. The somastatin analogue octreotide has been shown able to decrease morbidity after pancreatic resection. We studied the effect of perioperative treatment with octreotide on technical complications after pancreas transplant. PATIENTS AND METHODS: Seventeen recipients of bladder-drained transplant were randomized to receive either octreotide, 100 microg TID SQ for 5 days after transplant (n = 10) or no additional treatment (n = 7). We compared the two groups in terms of patient and graft survival and incidence of graft pancreatitis, intra- abdominal infections, and anastomotic leaks. RESULTS: In the untreated group, 1 patient developed a bladder leak and 2 had intra-abdominal infections, while no complications occurred in the octreotide-treated patients (P = 0.05). Six-month patient and pancreas survival was 100% and 90%, respectively, in octreotide-treated patients versus 86% and 86% in the control group (P = NS). CONCLUSION: Perioperative treatment with octreotide seems able to reduce the incidence of technical complications after pancreas transplantation.

KY: Adult; Amylases / blood / urine; Comparative-Study; Data-Interpretation-Statistical; Female; Follow-Up-Studies; Graft-Survival; Hormones-Synthetic /


Search
Clear
Print+Save
Setup
Handbook
Glossary
Help
Exit
About the Cochrane Collaboration (0 hits, 90 total)
The Cochrane Collaboration (0 hits, 1 total)
Collaborative Review Groups - CRGs (0 hits, 46 total)
Fields (0 hits, 10 total)
Methods Working Groups - MWGs (0 hits, 16 total)
Networks (0 hits, 1 total)
Centres (0 hits, 15 total)
Australasian Cochrane Centre
Baltimore Center
Brazilian Cochrane Centre
Canadian Cochrane Centre
Centre Cochrane Français
Dutch Cochrane Centre
Comments/Criticisms
Back
Find
Outline
Canadian Cochrane Centre
Contact address :
Kathie Clark
Coordinator
Canadian Cochrane Centre
Health Information Research Unit, McMaster University
1200 Main Street West
Hamilton
Ontario
Off-line

The Cochrane Collaboration Handbook

File Edit Bookmark Help

Contents | Search | Back | History | << | >>

The Cochrane Collaboration Handbook Version 3.0.2 (compiled September 1997)

Contents

About the Handbook
What's new?

1.INTRODUCTION
2. FORMAT OF A COCHRANE REVIEW
3. DEVELOPING A PROTOCOL
4. FORMULATING THE PROBLEM
5. LOCATING AND SELECTING STUDIES
6. CRITICAL APPRAISAL OF STUDIES
7. COLLECTING DATA
8. ANALYSING AND PRESENTING RESULTS
9. INTERPRETING RESULTS
10. IMPROVING AND UPDATING REVIEWS
11. REVIEWS USING INDIVIDUAL PATIENT DATA

Appendices
Glossary

Summary of analyses

MetaView: Tables and Figures

Discussion

Doppler ultrasound is a relatively new technique to be applied to the study of feto-placental and/or uteroplacental circulatory dynamics. It has been evaluated by more randomized trials than has any other biophysical test of fetal growth or wellbeing. However, one of the trials that was included in a previous version of this review, has now been excluded because of doubts about the veracity of the data (see Excluded Studies table). This is the London (KCH) trial conducted by JM Pearce.

Conclusions

Implications for practice

Doppler ultrasound is a relatively new technique to be applied to the study of feto-placental and/or uteroplacental circulatory dynamics. Although it has been evaluated more rigorously than any other biophysical test of fetal growth or wellbeing, the numbers of women included in the different trials fall somewhat short of the power calculations which preceded a recent systematic review of this subject (Alfirevic and Neilson 1995). The numbers have decreased with exclusion of the Pearce trial data (as has the confidence with which one can ascribe the reduced perinatal mortality rate to the use of Doppler ultrasound). Readers will either regard these findings as sufficiently sound to justify the use of Doppler ultrasound in high risk pregnancies (especially those complicated by hypertension or presumed fetal growth restriction), or see the findings as sufficiently promising to justify a larger, and hopefully definitive, trial.

Implications for research

As discussed, a case can be made for a larger trial of Doppler ultrasound than has been mounted hitherto. It is not clear if the study of uteroplacental arteries makes any real contribution or not.

MetaView 3.1 - [Perinatal deaths (Doppler in high risk pregnancies)]

File Display Sort Statistic Window Next Outcome Prev Outcome Help

Comparison: **Doppler ultrasound in high risk pregnancies**
Outcome: **Perinatal deaths**




Print
What to output
The current item only
Selected items
Which items to output
Only items marked with a tick (check)
All hits
How to output multiple items
Exactly as they appear in the index
Short document, short reference
Full document, complete reference
Print
Print Setup
Save
Cancel

Pentoxifylline in acute stroke.